

UR-1264

ER-40685-718

hep@xxx/9207014

A Note on “Anomalous” Chiral Gauge Theories

Zhu Yang

Department of Physics and Astronomy

University of Rochester

Rochester, NY 14627

bitnet:yang@uorhep

Abstract

We show that, the lattice regularization of chiral gauge theories proposed by Kaplan, when applied to a (2+1)-dimensional domain wall, produces a (1+1)-dimensional theory at low energy even if gauge anomaly produced by chiral fermions does not cancel. But the corresponding statement is not true in higher dimensions.

Chiral gauge theories deserve deeper study because nature exhibits left-right asymmetry. Recently Kaplan [1] has proposed a new lattice regularization of chiral gauge theories (see also [2]). The idea is to use the fermion zero modes trapped in a $2n+1$ dimensional domain wall as matter fields of $2n$ dimensions. Under suitable conditions, the fermion is chiral even on a lattice. The construction reproduces correct anomalous Ward identities. There is one catch, however: One has to make sure that the gauge fields live only in $2n$ dimensions as well. Kaplan argues that this can be true if and only if gauge currents in the $2n$ dimensional theory is anomaly free. Thus the method can be a very powerful tool to study non-perturbative aspects of chiral gauge theories. Some numerical results have already been obtained in $1+1$ dimension chiral Schwinger model [3].

On the other hand, $2n$ dimensional anomalous chiral gauge theories do not behave in the same way for all n . For example, an anomalous chiral gauge theory can be consistently quantized in $D=2$ at all energy scales, but not in $D=4$ or higher, see [4] for a review. Thus it would be disturbing if the Kaplan regularization did not comply with this fact, since it is so promising. In this note we show that it does. The difference between $D=2$ and $D > 2$ theories, in the present language, is due to the role of Chern-Simons terms in $D+1$ dimensions. So the Kaplan proposal gives a satisfactory lattice regularization of “anomalous” $1+1$ dimensional chiral gauge theory.

Since the problem arises in the continuum language as well, we will not use lattice cutoff explicitly. Following Kaplan, consider a fermion in $2n+1$ dimensions with a position dependent mass,

$$L = \bar{\psi}(\not{\partial} + m(s))\psi \tag{1}$$

where $m(s)$ satisfies is a step function with $m(0) = 0$. It can be shown that the fermion zero mode is chiral, due to the normalizability of the zero modes.

When coupled to some gauge field, the massive fermions produce a Goldstone-Wilczek[5]

current [6, 7, 8]

$$J_\mu \sim \text{sgn}m(s)\epsilon^{\alpha_1\beta_1\ldots\alpha_n\beta_n}F_{\alpha_1\beta_1}\ldots F_{\alpha_n\beta_n}. \quad (2)$$

This current is not divergenceless, because of the sign function, but it is cancelled by contribution from the fermion zero modes trapped on the wall[9]. This is in accord with the general expectation that there is no anomaly of continuous symmetry in odd dimensions.

Superficially it appears that one has successfully devised a 2n dimensional chiral gauge theory, or at least a model of chiral fermions coupled to gauge fields. One must, as Kaplan pointed out, be careful about the gauge field dynamics, making sure it is correct in 2n dimensional sense. Kaplan argues that for a continuum Yang-Mills action

$$L_{YM} = \beta_1 F_{ij}^2 + \beta_2 F_{is}^2, \quad (3)$$

the equation of motion for A_μ is

$$\begin{aligned} \beta_1 D_j F_{ij} + D_s \beta_2 F_{is} &= J_i, \\ \beta_2 D_i F_{si} &= J_s. \end{aligned} \quad (4)$$

In general J_s does not vanish, so one cannot set β_2 to zero. One is not studying 2n dimensional gauge theory.

However for the spacial case of n=1, the long wave length of the gauge fields off the wall can be derived effectively from the following action

$$L = \beta_1 F_{ij}^2 + \beta_2 F_{is}^2 + \frac{\text{sgn}m(s)}{4\pi} (AdA + \frac{2}{3}A^3). \quad (5)$$

This is more or less the same as the topologically massive gauge theory [10, 6], except for some position dependent coupling constants. It simply implies that there is nothing in low-energy physics in 2+1 dimensions, since the gauge bosons are massive. More correctly, it is a Chern-Simons topological field theory, which is known to possess no local propagating degrees of freedom [11]. The scale at which nontrivial dynamics sets in is given by $1/\beta_2$.

On the other hand the typical energy scale in 2 dimensions is $\sqrt{m/\beta_2}$. The latter can be much lower than the former if $m \ll \beta$. So a generically non-zero β_2 will still have no effect on the low-energy dynamics if $\beta_2(0) = 0$ and otherwise β_1 , and β_1 is tuned to be much smaller than $1/m$. In this particular case, the model does look like a 2 dimensional theory. The theory could be termed “anomalous”, because the chiral fermions do produce anomaly. But it is cancelled because of the massive fermion contribution from the extra dimension. For $n \geq 2$, the resulting Chern-Simons term has no low energy effect, so that the gauge bosons are massless in $2n + 1$ dimensions, and the regularization fails to produce a truly $2n$ dimensional theory. Note even if the gauge anomaly cancels, one must still tune the parameters to cancel induced F_{si}^2 term from matter fields.

We put quotation marks on the word anomalous because, as is well-known, a 2D “anomalous” gauge theory is not really anomalous, because the gauge degrees of freedom happen to be described by a chiral Lagrangian as well, and the resulting model is renormalizable and can even be finite. In our case, the extra degrees of freedom are obviously related to the Chern-Simons gauge theory with a boundary [11, 12], which is a chiral WZW theory by itself. In fact, the gapless edge excitation of a quantum Hall system is a physical realization of 2D chiral gauge theory [13].

In conclusion, we have shown that Kaplan’s regularization works for “anomalous” gauge theories in D=2 as well. Our observation further supports the viability of that proposal.

Acknowledgement

This work is supported in part by U.S. Department of Energy Contract No. DE-AC02-76ER13065.

References

- [1] D. Kaplan, San Diego preprint UCSD/PTH 92-16.
- [2] D. Boyanovsky, E. Dagotto and E. Fradkin, Nucl. Phys. **B285** (1987) 340; E. Dagotto, E. Fradkin and A. Moreo, Phys. Lett. **B172** (1986) 383.
- [3] K. Jansen, San Diego preprint UCSD/PTH 92-18.
- [4] J. Preskill, Ann. Phys. **210** (1991) 323.
- [5] J. Goldstone and F. Wilczek, Phys. Rev. Lett. **47** (1981) 986.
- [6] S. Deser, R. Jackiw and S. Templeton, Phys. Rev. Lett. **48** (1982) 975; Ann. Phys. (N.Y.) **140** (1982) 372.
- [7] A.N. Redlich, Phys. Rev. Lett. **52** (1984) 18; Phys. Rev. **D29** (1984) 2366.
- [8] S. Naculich, Nucl. Phys. **B296** (1988) 837.
- [9] C.G. Callan and J.A. Harvey, Nucl. Phys. **B250** (1985) 427.
- [10] J. Schonfeld, Nucl. Phys. **B185** (1981) 157.
- [11] E. Witten, Comm. Math. Phys. **121** (1989) 351.
- [12] G. Moore and N. Seiberg, Phys. Lett. **B220** (1989) 422.
- [13] X.G. Wen, preprint NSF-ITP-89-157; M. Stone, Ann. Phys. (N.Y.) **207** (1991) 38.